

High Energy Resummation of Drell-Yan Processes

Simone Marzani^a, Richard D. Ball^b,

^a *School of Physics and Astronomy, University of Manchester,
Oxford Road, Manchester M13 9PL, England, UK*

^b *School of Physics, University of Edinburgh,
Mayfield Rd, Edinburgh EH9 3JZ, Scotland, UK*

Abstract

We present a computation of the inclusive Drell-Yan production cross-section in perturbative QCD to all orders in the limit of high partonic centre-of-mass energy. We compare our results to the fixed order NLO and NNLO results in $\overline{\text{MS}}$ scheme, and provide predictions at N³LO and beyond. Our expressions may be used to obtain fully resummed results for the inclusive cross-section.

December 2008

1 Introduction

Accurate perturbative calculations of benchmark inclusive cross-sections are an essential component of the LHC discovery programme. The most important of these benchmark processes are the Drell-Yan processes: production of $\mu^+\mu^-$ pairs, and the closely related vector boson production processes. Currently these cross-sections are known at NNLO in perturbative QCD [1–4]. The resummation of threshold logarithms is known up to N³LL [5–8].

When the invariant mass Q of the particles produced in the final state is well below the centre-of-mass energy, the typical values of x of the colliding partons may be rather small: $x_1x_2 = Q^2/S \ll 1$. In fact this is true of most LHC processes: only when producing very massive states close to threshold do both partons carry a large fraction of the incoming longitudinal momentum. Whenever x is small, logarithms of x may spoil the perturbation series. Thus accurate calculations require the computation of the coefficients of these logarithms, and if they are large it may be necessary to resum them.

The resummation of small- x logarithms in the perturbative evolution of parton distribution functions at NLL is by now well understood (see for example ref. [9] for a recent review). The general procedure for resumming hard cross-sections at the same order through k_T -factorization is known [10, 11], and its implementation when the coupling runs understood [12]. Calculations have been performed for photoproduction processes [10, 12], deep inelastic processes [11, 13], hadroproduction of heavy quarks [10, 12, 14, 15], and gluonic Higgs production both in the pointlike limit [16], and for finite top mass [17, 18]. However for Drell-Yan and vector boson production the resummed hard cross-section has yet to be computed. It is the purpose of this paper to perform this calculation, determining the coefficients of the leading high energy singularities of the Drell-Yan coefficient function to all orders in perturbation theory. This will enable the accurate evaluation of small- x corrections to these benchmark processes, soon to be measured at LHC.

2 The Drell-Yan cross-section

We wish to study the high energy behaviour of the Drell-Yan cross-section. We consider n_f quarks q_i with electric charge e_i . The cross-section for the production of a lepton pair via an off-shell photon with squared momentum $q^2 = Q^2$ can then be written as:

$$\sigma(\tau_h, Q^2) = \sigma_0(Q^2) \sum_{a,b=q_i,\bar{q}_j,g} \int_{\rho}^1 \frac{dx_1}{x_1} \int_{\rho}^1 \frac{dx_2}{x_2} D_{ab}(\frac{\tau_h}{x_1x_2}; \alpha_s(Q^2)) F_a(x_1, Q^2) F_b(x_2, Q^2), \quad (1)$$

where $\tau_h = Q^2/S$, and $F_a(x, Q^2) = x f_a(x, Q^2)$ is the integrated parton density for parton a , the indices a and b running over the different initial partons $q_i, \bar{q}_j, g$. The LO partonic cross-section is simply

$$\sigma_0(Q^2) = \frac{\alpha}{Q^2} \frac{4\pi}{3N_c} \langle e^2 \rangle, \quad (2)$$

where $\langle e^2 \rangle \equiv \frac{1}{n_f} \sum_i e_i^2$: the dimensionless coefficient functions D_{ab} then contain all the QCD radiative corrections.




Figure 1: (a) LO and (b) NLO diagrams contributing to $D_{q\bar{q}}$, and (c) a NNLO diagram contributing to the flavour disconnected piece $\tilde{D}_{q\bar{q}}$. Both quark and antiquark must carry the same flavour.

In what follows we will employ a Mellin transform in τ_h to undo the convolutions in eqn. (1): defining

$$\sigma(N, Q^2) = \int_0^1 d\tau_h \tau_h^{N-1} \sigma(\tau_h, Q^2), \quad (3)$$

the factorization eqn. (1) becomes simply

$$\sigma(N, Q^2) = \sigma_0(Q^2) \sum_{a,b=q_i,\bar{q}_j,g} D_{ab}(N; \alpha_s(Q^2)) F_a(N, Q^2) F_b(N, Q^2). \quad (4)$$

Clearly $D_{ab} = D_{ba}$, and thus since the strong interaction is CP even $D_{q_i q_j} = D_{\bar{q}_i \bar{q}_j}$, $D_{q_i \bar{q}_j} = D_{\bar{q}_i q_j}$ and $D_{q_i g} = D_{\bar{q}_i g}$. Thus if we define singlet and nonsinglet quark (plus antiquark) distributions and quark-antiquark luminosities as

$$\begin{aligned} Q &\equiv \sum_i (Q_i + \bar{Q}_i), & \langle Q \rangle &\equiv \frac{1}{\langle e^2 \rangle} \sum_i e_i^2 (Q_i + \bar{Q}_i) \\ \{Q\bar{Q}\} &\equiv \sum_i Q_i \bar{Q}_i & \langle Q\bar{Q} \rangle &\equiv \frac{1}{\langle e^2 \rangle} \sum_i e_i^2 Q_i \bar{Q}_i, \end{aligned} \quad (5)$$

where $Q_i = xq_i$ etc, then we can decompose the factorization eqn. (4) as

$$\begin{aligned} \sigma &= \sigma_0 [D_{q\bar{q}} \langle Q\bar{Q} \rangle + D_{qq} \langle Q \rangle Q + D_{qg} \langle Q \rangle G \\ &\quad + n_f \tilde{D}_{q\bar{q}} \{Q\bar{Q}\} + n_f \tilde{D}_{qq} QQ + 2n_f \tilde{D}_{qg} QG + n_f \tilde{D}_{gg} GG]. \end{aligned} \quad (6)$$

Inevitably this flavour decomposition is rather more complicated than in DIS (see ref. [11]) because here one must consider the flavours of three partons: the two incoming ones and the one from which the virtual photon is emitted.

We now consider the behaviour of the perturbation series for the coefficients $D_{ab}(N; \alpha_s(Q^2))$. At LO the only process is $q\bar{q} \rightarrow \gamma^*$ and thus $D_{q\bar{q}} = 1$ (fig.1a), while all the rest are zero. At NLO the channel $gg \rightarrow q\gamma^*$ opens: $D_{qg} = \mathcal{O}(\alpha_s)$ (fig.2). Processes with two gluons, two quarks, or two antiquarks in




Figure 2: NLO diagrams contributing to D_{qg} .




Figure 3: NNLO diagrams contributing to $\tilde{D}_{qg}$ at NNLO.

the initial state can only occur at NNLO: $\tilde{D}_{gg} = \mathcal{O}(\alpha_s^2)$ (fig.3), and $D_{qq} = \mathcal{O}(\alpha_s^2)$ (fig.5a and 5b). At this order there is also a “flavour disconnected” process contributing to $q\bar{q} \rightarrow \gamma^*$: $\tilde{D}_{q\bar{q}} = \mathcal{O}(\alpha_s^2)$ (fig.1c). Flavour disconnected processes contribute also to $qg \rightarrow X\gamma^*$ ($\tilde{D}_{qg} = \mathcal{O}(\alpha_s^3)$, fig.4c) and $qq \rightarrow X\gamma^*$ ($\tilde{D}_{qq} = \mathcal{O}(\alpha_s^4)$, fig.5c): note that these each have a $gg \rightarrow q\bar{q}\gamma^*$ subprocess. It is easy to see that all these flavour disconnected pieces are two powers of α_s down compared to the corresponding flavour connected piece, and that they must be proportional to n_f .

Now consider the behaviour of the coefficient functions at high energy, that is as $N \rightarrow 0$. Only some of the higher order corrections contain high energy logarithms, and these are the ones we wish to isolate and compute. To see how these logarithms arise, and thus deduce the general pattern, note that at LO the gg anomalous is singular, $\gamma_{gg} \sim \frac{\alpha_s}{N}$, while the qq anomalous dimension is not, $\gamma_{qq} \sim \alpha_s$. Thus the NLO corrections to $D_{q\bar{q}}$ due to gluon emission from a quark line (fig.1b) are $\mathcal{O}(\alpha_s)$: further gluon emissions from quark lines give contributions of $\mathcal{O}(\alpha_s^k)$. Similarly the flavour disconnected piece $\tilde{D}_{q\bar{q}}$ has no high energy logarithms. In fact both the terms in eqn. (6) in which an incoming quark and antiquark annihilate to produce the vector boson are completely regular as $N \rightarrow 0$, in the same way that the nonsinglet contribution to DIS structure functions is regular:

$$D_{q\bar{q}} = 1 + C_F \left(\frac{4}{3} \pi^2 - \frac{7}{2} \right) \frac{\alpha_s}{2\pi} + \mathcal{O}(\alpha_s^2), \quad \tilde{D}_{q\bar{q}} = \mathcal{O}(\alpha_s^2). \quad (7)$$




Figure 4: (a) and (b) are NNLO diagrams contributing to D_{gg} , which introduce high energy logarithms, while (c) is a N³LO diagram contributing to the flavour disconnected piece $\tilde{D}_{gg}$.




Figure 5: (a) and (b) are NNLO diagrams contributing to D_{qq} , while (c) is an N⁴LO diagram contributing to the flavour disconnected piece $\tilde{D}_{qq}$.

However the NNLO contribution to D_{gg} can be singular, due to gluon emission from the incoming gluon line (fig.4a and 4b), and is thus $\mathcal{O}(\alpha_s \frac{\alpha_s}{N})$. Further gluon emissions from the incoming gluon give extra powers of $\frac{\alpha_s}{N}$, while gluon emissions from a quark line just give an extra α_s , so the most singular contributions are $\mathcal{O}(\alpha_s (\frac{\alpha_s}{N})^k)$, with $k \geq 0$, i.e. NLL x . Similarly the N³LO contribution to D_{gg} is $\mathcal{O}(\alpha_s^2 \frac{\alpha_s}{N})$, due to gluon emission from one of the incoming gluons: in general the most singular contributions here are $\mathcal{O}(\alpha_s^2 (\frac{\alpha_s}{N})^k)$, $k \geq 0$ i.e. NNLL x . Finally consider D_{qq} : at NNLO besides gluon emission from quark lines, one of the initial quarks can turn into a gluon, the rest of the process then proceeding as for D_{gg} (fig.5a and 5b). Since $\gamma_{gq} \sim \frac{C_F}{C_A} \frac{\alpha_s}{N}$, these graphs are also singular, $\mathcal{O}(\alpha_s \frac{\alpha_s}{N})$. Further emissions from the gluon line then give contributions of $\mathcal{O}(\alpha_s (\frac{\alpha_s}{N})^k)$, $k \geq 1$, and are thus again NLL x . In fact these graphs are at high energy identical to the corresponding graphs for D_{gg} , save for the extra factor of $\frac{C_F}{C_A}$ due to the incoming quark rather than incoming gluon, so there is a colour-charge relation [11] between the leading singularities of D_{qq} and those of D_{gg} . Similar considerations apply to the disconnected contributions $\tilde{D}_{qq}$ and $\tilde{D}_{gg}$: the most singular contributions to $\tilde{D}_{gg}$ are $\mathcal{O}(\alpha_s^2 (\frac{\alpha_s}{N})^k)$, $k \geq 1$, and related through a factor of $\frac{C_F}{C_A}$ to those of D_{gg} (see fig. 4c), while those of $\tilde{D}_{qq}$ are $\mathcal{O}(\alpha_s^2 (\frac{\alpha_s}{N})^k)$,

$k \geq 2$, and related through a factor of $(\frac{C_F}{C_A})^2$ to those of D_{gg} (see fig. 5c).

Summarising, in the high energy limit the leading singularities of the coefficient functions have the schematic structure

$$\begin{aligned}
D_{qq}(N, \alpha_s) &= \frac{C_F}{C_A} \sum_{k=1}^{\infty} \mathcal{O}(\alpha_s (\frac{\alpha_s}{N})^k), \\
D_{qg}(N, \alpha_s) &= \mathcal{O}(\alpha_s) + \sum_{k=1}^{\infty} \mathcal{O}(\alpha_s (\frac{\alpha_s}{N})^k), \\
\tilde{D}_{qq}(N, \alpha_s) &= (\frac{C_F}{C_A})^2 \sum_{k=2}^{\infty} \mathcal{O}(\alpha_s^2 (\frac{\alpha_s}{N})^k), \\
\tilde{D}_{qg}(N, \alpha_s) &= \frac{C_F}{C_A} \sum_{k=1}^{\infty} \mathcal{O}(\alpha_s^2 (\frac{\alpha_s}{N})^k), \\
\tilde{D}_{gg}(N, \alpha_s) &= \mathcal{O}(\alpha_s^2) + \sum_{k=1}^{\infty} \mathcal{O}(\alpha_s^2 (\frac{\alpha_s}{N})^k).
\end{aligned} \tag{8}$$

It follows that while the quark-quark and quark-gluon coefficient functions are NLL x , the gluon-gluon coefficient function and the disconnected contributions to quark-quark and quark-gluon are all NNLL x , and thus need not be considered in a NLL x calculation (though they may still be numerically large, since the gluon distribution is rather larger than the quark distribution). Our strategy in the rest of this paper will thus be to use k_T -factorization to compute the leading high energy singularities of the D_{qg} coefficient function to all order in perturbation theory, and then deduce those of D_{qq} using colour-charge relations. The high energy singularities are found by considering the off-shell process $g^* q \rightarrow \gamma^* q$ calculated from the graphs in fig. 2 with the incoming gluon off-shell, in just the same way as one determines the coefficient function C_{2g} in DIS using the off-shell process $\gamma^* g^* \rightarrow q\bar{q}$ [11]. The computation is complicated by the fact that just as in the DIS calculation there are collinear singularities as well as high energy singularities, so one has to take care to factorize the collinear singularities before extracting the high energy ones. In the next section we explain in some detail how this can be done.

The hadroproduction of vector bosons, either $W^\pm$ or Z , has a very similar structure to Drell-Yan: in Mellin space the cross section can be written as

$$\sigma^V(N, m_V^2) = \sigma_0^V \sum_{a,b=q,\bar{q},g} D_{ab}^V(N; \alpha_s(Q^2)) F_a(N, Q^2) F_b(N, Q^2). \tag{9}$$

where now

$$\sigma_0^V = \frac{\pi}{N_c} \sqrt{2} G_F \tag{10}$$

and G_F is the Fermi constant. The decomposition eqn. (4) is the same up to threshold effects, except that for Z -production e_i^2 in eqn. (5) is replaced by $v_i^2 + a_i^2$, where v_i, a_i are the vector and axial couplings of the Z bosons to the quarks, while for $W^\pm$ production

$$\langle Q \rangle = Q, \quad \langle Q\bar{Q} \rangle = \sum_{ij} |V_{ij}|^2 Q_i \bar{Q}_j, \tag{11}$$

where V_{ij} is the CKM matrix, $\sum_j V_{ij}V_{jk}^* = \delta_{ik}$, so the disconnected pieces of the quark-quark and quark-gluon coefficient functions may be simply added to the connected pieces. The perturbative QCD corrections to vector boson production are the same as the Drell-Yan ones at NLO, and they only begin to differ at NNLO because of the diagram with two incoming gluons and an internal quark loop, which gives an $\mathcal{O}(\alpha_s^2)$ contribution to the Z boson cross-section but vanishes in the case of a virtual photon or $W^\pm$. The NLL x singularities are thus determined by the off-shell processes $g^*q \rightarrow W^\pm q'$ and $g^*q \rightarrow Zq$ at $\mathcal{O}(\alpha_s)$ and are the same as those for Drell-Yan: the NNLL x singularities for Z production would however receive an extra contribution.

3 High energy factorization

The general structure of high energy singularities outlined in the previous section arises from the k_T -factorization formalism [10], and in particular its extension by Catani and Hautmann to deal with situations in which the hard cross-section contains collinear singularities which must be factorized consistently [11]. In this section we explain how this works for Drell-Yan processes.

We saw in the previous section how the singular contributions to the cross-section eqn. (6) are of essentially two kinds: those in which the electroweak boson is emitted directly from one of the incoming quarks or antiquarks, and those (the ‘disconnected’ pieces) where it is not. While the former can only acquire high energy logarithms through gluon emission from the second parton, the former acquire high energy logarithms through gluon emission from both partons. The dimensional regularised form of k_T -factorization for the singular part of the Drell-Yan cross-section is thus (in dimensional regularization)

$$\begin{aligned} \sigma_s(N, Q^2) = & \sigma_0(Q^2) \left\{ \int \frac{d^{2-2\varepsilon} \mathbf{k}}{\pi \mathbf{k}^2} \Sigma_{qg}(N, \frac{\mathbf{k}^2}{Q^2}; \alpha_s, \varepsilon) \langle Q^{(0)}(N, \mu, \varepsilon) \right. \\ & \left. (\mathcal{F}_q^{(0)}(N, \frac{\mathbf{k}^2}{\mu^2}; \alpha_s, \varepsilon) Q^{(0)}(N, \mu, \varepsilon) + \mathcal{F}_g^{(0)}(N, \frac{\mathbf{k}^2}{\mu^2}; \alpha_s, \varepsilon) G^{(0)}(N, \mu, \varepsilon)) \right. \\ & + n_f \int \frac{d^{2-2\varepsilon} \mathbf{k}_1}{\pi \mathbf{k}_1^2} \frac{d^{2-2\varepsilon} \mathbf{k}_2}{\pi \mathbf{k}_2^2} \Sigma_{gg}(N, \frac{\mathbf{k}_1^2}{Q^2}, \frac{\mathbf{k}_2^2}{Q^2}; \alpha_s, \varepsilon) \\ & \left. (\mathcal{F}_q^{(0)}(N, \frac{\mathbf{k}_1^2}{\mu^2}; \alpha_s, \varepsilon) Q^{(0)}(N, \mu, \varepsilon) + \mathcal{F}_g^{(0)}(N, \frac{\mathbf{k}_1^2}{\mu^2}; \alpha_s, \varepsilon) G^{(0)}(N, \mu, \varepsilon)) \right. \\ & \left. (\mathcal{F}_q^{(0)}(N, \frac{\mathbf{k}_2^2}{\mu^2}; \alpha_s, \varepsilon) Q^{(0)}(N, \mu, \varepsilon) + \mathcal{F}_g^{(0)}(N, \frac{\mathbf{k}_2^2}{\mu^2}; \alpha_s, \varepsilon) G^{(0)}(N, \mu, \varepsilon)) \right\}, \end{aligned} \quad (12)$$

where $\Sigma_{qg}(N, \frac{\mathbf{k}^2}{Q^2}; \alpha_s, \varepsilon)$ is the partonic cross-section for $qg \rightarrow \gamma^* q$ with the incoming gluon off-shell (given at leading order by the diagrams fig. 2, and thus $\mathcal{O}(\alpha_s)$), $\Sigma_{gg}(N, \frac{\mathbf{k}_1^2}{Q^2}, \frac{\mathbf{k}_2^2}{Q^2}; \alpha_s, \varepsilon)$ is the partonic cross-section for $gg \rightarrow \gamma^* q \bar{q}$ with both the incoming gluons off-shell (given at leading order by the diagrams fig. 3, and thus $\mathcal{O}(\alpha_s^2)$), $\mathcal{F}_q^{(0)}$ and $\mathcal{F}_g^{(0)}$ are the bare quark and gluon Green’s functions, and $Q^{(0)}$ and $G^{(0)}$ the bare integrated quark (plus antiquark) and gluon densities.¹ At LL x the quark

¹Note that our notation is slightly different to that in [11], since in particular we prefer to deal throughout with integrated parton densities.

Green's function is

$$\mathcal{F}_q^{(0)}(N, \frac{\mathbf{k}^2}{\mu^2}; \alpha_s, \varepsilon) = \frac{C_F}{C_A} \left[\mathcal{F}_g^{(0)}(N, \frac{\mathbf{k}^2}{\mu^2}; \alpha_s, \varepsilon) - \pi \mathbf{k}^2 \delta^{(2-2\varepsilon)}(\mathbf{k}) \right], \quad (13)$$

while the gluon Green's function is

$$\mathcal{F}_g^{(0)}(N, \frac{\mathbf{k}^2}{\mu^2}; \alpha_s, \varepsilon) = \gamma_{gg} R(\gamma_{gg}) \Gamma_{gg}(N; \alpha_s, \varepsilon) \left(\frac{\mathbf{k}^2}{\mu^2} \right)^{\gamma_{gg}}. \quad (14)$$

The gluon-gluon transition function (containing the collinear poles) is given by

$$\Gamma_{gg}(N; \alpha_s, \varepsilon) = \exp \left(\frac{1}{\varepsilon} \int_0^{\alpha_s S_\varepsilon} \frac{d\alpha}{\alpha} \gamma_{gg}(N, \alpha) \right), \quad (15)$$

where $S_\varepsilon = \exp[-\varepsilon(-\gamma_E + \ln 4\pi)]$ and $R(\gamma_{gg})$ is a scheme-dependent normalization factor, regular as $\varepsilon \rightarrow 0$: in the $\overline{\text{MS}}$ scheme

$$R(\gamma_{gg}) = \left\{ \frac{\Gamma(1 - \gamma_{gg}) \chi_0(\gamma_{gg})}{\Gamma(1 + \gamma_{gg}) [-\gamma_{gg} \chi'_0(\gamma_{gg})]} \right\}^{1/2} \exp \left\{ \gamma_{gg} \psi(1) + \int_0^{\gamma_{gg}} d\gamma \frac{\psi'(1) - \psi'(1 - \gamma)}{\chi_0(\gamma)} \right\}. \quad (16)$$

In the high energy limit

$$\gamma_{gg}(N, \alpha_s) = \gamma_s \left(\frac{\alpha_s}{N} \right) + \mathcal{O} \left(\alpha_s \left(\frac{\alpha_s}{N} \right)^k \right). \quad (17)$$

where $\gamma_s \left(\frac{\alpha_s}{N} \right) = \frac{\alpha_s}{N} + \mathcal{O} \left(\left(\frac{\alpha_s}{N} \right)^4 \right)$ is the naive dual to the BFKL kernel χ_0 .

We can use the k_T -factorization eqn. (12) to extract formulae for the bare coefficients in the expression eqn. (6) at high energy in terms of the off-shell partonic cross-sections Σ_{qg} and Σ_{gg} . For instance, comparing the coefficients of $\langle Q^{(0)} \rangle G^{(0)}$, we have

$$\begin{aligned} D_{qg}^{(0)}(N, Q^2; \alpha_s, \varepsilon) &= \int \frac{d^{2-2\varepsilon} \mathbf{k}}{\pi \mathbf{k}^2} \Sigma_{qg}(N, \frac{\mathbf{k}^2}{Q^2}; \alpha_s, \varepsilon) \mathcal{F}_g^{(0)}(N, \frac{\mathbf{k}^2}{\mu^2}; \alpha_s, \varepsilon) \\ &= \int \frac{d^{2-2\varepsilon} \mathbf{k}}{\pi \mathbf{k}^2} \Sigma_{qg}(N, \frac{\mathbf{k}^2}{Q^2}; \alpha_s, \varepsilon) \left(\frac{\mathbf{k}^2}{\mu^2} \right)^{\gamma_{gg}} \gamma_{gg} R(\gamma_{gg}) \Gamma_{gg}(N; \alpha_s, \varepsilon), \end{aligned} \quad (18)$$

where in the second line we substituted the gluon Green's function (14). Introducing $\xi = \mathbf{k}^2/Q^2$, and defining the impact factor

$$h_{qg}(N, M; \alpha_s) \equiv M^2 \int_0^\infty d\xi \xi^{M-1} \Sigma_{qg}(N, \xi; \alpha_s, 0) \quad (19)$$

we can write this as

$$D_{qg}^{(0)}(N, Q^2; \alpha_s, \varepsilon) = \left(\frac{Q^2}{\mu^2} \right)^{\gamma_{gg}} h_{qg}(N, \gamma_{gg}; \alpha_s) \gamma_{gg}^{-1} R(\gamma_{gg}) \Gamma_{gg}(N; \alpha_s, \varepsilon) + \mathcal{O}(\alpha_s^2 \left(\frac{\alpha_s}{N} \right)^k). \quad (20)$$

Note that we can set $\varepsilon = 0$ in the impact factor, since collinear singularities are now regularised by $M \neq 0$. The reason for the extra factor of M^2 in eqn. (19) will become clear shortly.

Similarly, comparing coefficients of $\langle Q^{(0)} \rangle Q^{(0)}$ we find

$$\begin{aligned}
D_{qq}^{(0)}(N, Q^2; \alpha_s, \varepsilon) &= \frac{C_F}{C_A} \left[\left(\frac{Q^2}{\mu^2} \right)^{\gamma_{gg}} h_{qq}(N, \gamma_{gg}; \alpha_s) \gamma_{gg} R(\gamma_{gg}) \Gamma_{gg}(N; \alpha_s, \varepsilon) \right. \\
&\quad \left. - \Sigma_{qq}(N, 0; \alpha_s, \varepsilon) \right] + \mathcal{O}(\alpha_s^2 (\frac{\alpha_s}{N})^k), \\
&= \frac{C_F}{C_A} \left[D_{qq}^{(0)}(N, Q^2; \alpha_s, \varepsilon) - \Sigma_{qq}(N, 0; \alpha_s, \varepsilon) \right] + \mathcal{O}(\alpha_s^2 (\frac{\alpha_s}{N})^k), \tag{21}
\end{aligned}$$

the second term (which is also singular) coming from the delta-function subtraction in eqn. (13). Similar expressions for the bare coefficients $\tilde{D}_{qq}$, $\tilde{D}_{qg}$ and $\tilde{D}_{gg}$ may be extracted in terms of double Mellin transforms of Σ_{gg} : we will ignore these in what follows as they are $\mathcal{O}(\alpha_s^2)$.

4 Collinear factorization at high energy

We now need to factorize the collinear singularities in order to absorb them into renormalized parton distribution functions. In the case of a collinear safe process such as heavy flavour hadroproduction [14, 15] this is relatively straightforward: all the ε poles are in the Green's functions, and in the $\overline{\text{MS}}$ factorization scheme are absorbed into the bare gluon distribution, i.e. $G = \Gamma_{gg} G^{(0)}$. However when we consider a process in which the partonic cross-section is not collinear safe, such as DIS or DY, the factorization of the collinear singularities is rather more involved [11].

Firstly, we must separate out the valence and nonsinglet quark distributions from the singlet, since the former factorize multiplicatively while the latter mixes with the gluon. Defining

$$V_i \equiv Q_i - \bar{Q}_i, \quad N_i \equiv Q_i + \bar{Q}_i - \frac{1}{n_f} Q \tag{22}$$

so $\sum_i N_i = 0$, we can rewrite the various quantities (5) appearing in eqn. (6)

$$\begin{aligned}
\langle Q \rangle &= \frac{1}{\langle e^2 \rangle} \sum_i e_i^2 (N_i + \frac{1}{n_f} Q) = \langle N \rangle + Q, \\
\{Q\bar{Q}\} &= \frac{1}{4} \sum_i ((N_i + \frac{1}{n_f} Q)^2 - V_i^2) = \frac{1}{4} \{N^2\} - \frac{1}{4} \{V^2\} + \frac{1}{4n_f} Q^2, \\
\langle Q\bar{Q} \rangle &= \frac{1}{4} \frac{1}{\langle e^2 \rangle} \sum_i e_i^2 ((N_i + \frac{1}{n_f} Q)^2 - V_i^2) = \frac{1}{4} \langle N^2 \rangle - \frac{1}{4} \langle V^2 \rangle + \frac{1}{2n_f} \langle N \rangle Q + \frac{1}{4n_f} Q^2.
\end{aligned} \tag{23}$$

When written in terms of N , V , Q and G the Drell-Yan cross-section eqn. (6) separates into three distinct pieces: $\sigma = \sigma^{NS} + \sigma^S + \sigma^{SS}$, where

$$\sigma^{NS} = \sigma_0 \left[\frac{1}{4} D_{q\bar{q}} (\langle N^2 \rangle - \langle V^2 \rangle) + \frac{1}{4} n_f \tilde{D}_{q\bar{q}} (\{N^2\} - \{V^2\}) \right], \tag{24}$$

$$\sigma^S = \sigma_0 \left[(D_{qq} + \frac{1}{2n_f} D_{q\bar{q}}) \langle N \rangle Q + D_{qg} \langle N \rangle G \right], \tag{25}$$

$$\begin{aligned}
\sigma^{SS} &= \sigma_0 \left[(D_{qq} + \frac{1}{4n_f} D_{q\bar{q}}) Q Q + D_{qg} Q G \right. \\
&\quad \left. + (n_f \tilde{D}_{qq} + \frac{1}{4n_f} \tilde{D}_{q\bar{q}}) Q Q + 2n_f \tilde{D}_{qg} Q G + n_f D_{gg} G G \right]. \tag{26}
\end{aligned}$$

Now the factorization of collinear singularities for the valence and nonsinglet distributions V_i and N_i is multiplicative:

$$V_i(N, \alpha_s) = \Gamma_{qq}^V(N, \alpha_s, \varepsilon) V_i^{(0)}(N, \alpha_s, \varepsilon), \quad N_i(N, \alpha_s) = \Gamma_{qq}^N(N, \alpha_s, \varepsilon) N_i^{(0)}(N, \alpha_s, \varepsilon) \quad (27)$$

so the nonsinglet cross-section σ^{NS} factorizes multiplicatively. Furthermore the transition functions Γ_{qq}^V and Γ_{qq}^N contain no high energy singularities: at high energy the both reduce to unity up to corrections of $\mathcal{O}(\alpha_s)$. However singlet quark Q and gluon G mix:

$$Q = \Gamma_{qq} Q^{(0)} + 2n_f \Gamma_{qg} G^{(0)} \quad G = \Gamma_{gq} Q^{(0)} + \Gamma_{gg} G^{(0)}, \quad (28)$$

and the transition functions now contain high energy singularities. These were determined to NLL x in [11]: for example

$$\Gamma_{qg}(N, \alpha_s, \varepsilon) = \frac{1}{\varepsilon} \int_0^{\alpha_s S_\varepsilon} \frac{d\alpha}{\alpha} \gamma_{qg}(N, \alpha) \Gamma_{gg}(N, \alpha, \varepsilon), \quad (29)$$

where Γ_{gg} is given by eqn. (15), and thus Γ_{qg} contains (singular) terms of $\mathcal{O}(\alpha_s(\frac{\alpha_s}{N})^k)$. Γ_{qg} and Γ_{gg} are given in terms of Γ_{gg} and Γ_{qg} by colour-charge relations:

$$\Gamma_{gg}(N, \alpha_s, \varepsilon) = \frac{C_F}{C_A} [\Gamma_{gg}(N, \alpha_s, \varepsilon) - 1] \quad (30)$$

while writing $\Gamma_{qq} = 1 + 2n_f \Gamma_{qq}^{PS}$,

$$\Gamma_{qq}^{PS}(N, \alpha_s, \varepsilon) = \frac{1}{\varepsilon} \int_0^{\alpha_s S_\varepsilon} \frac{d\alpha}{\alpha} [\gamma_{qq} + \gamma_{qg} \Gamma_{gg}] = \frac{C_F}{C_A} [\Gamma_{qq}(N, \alpha_s, \varepsilon) - \frac{\alpha_s}{\varepsilon} \gamma_{qq}^0(N)], \quad (31)$$

since at high energy $\gamma_{qq}(N, \alpha_s) = \frac{C_F}{C_A} [\gamma_{qq}(N, \alpha_s) - \alpha_s \gamma_{qq}^0(0)]$, where $\gamma_{qq}(N, \alpha_s) = \alpha_s \gamma_{qq}^0(N) + \mathcal{O}(\alpha_s^2)$.

Thus if we consider the collinear factorization of the singlet cross-section σ^S , substituting eqns. (27,28) into eqn. (25), the gluon and quark contributions to σ^S are given respectively by

$$\sigma_0^{-1} \sigma_g^S \equiv D_{qg}^{(0)} \langle N^{(0)} \rangle G^{(0)} = \Gamma_{qq}^N \left[(D_{qq} + \frac{1}{2n_f} D_{q\bar{q}}) 2n_f \Gamma_{qg} + D_{qg} \Gamma_{gg} \right] \langle N^{(0)} \rangle G^{(0)}, \quad (32)$$

$$\sigma_0^{-1} \sigma_q^S \equiv (D_{qq}^{(0)} + \frac{1}{2n_f} D_{q\bar{q}}^{(0)}) \langle N^{(0)} \rangle Q^{(0)} = \Gamma_{qq}^N \left[(D_{qq} + \frac{1}{2n_f} D_{q\bar{q}}) \Gamma_{qq} + D_{qg} \Gamma_{gg} \right] \langle N^{(0)} \rangle Q^{(0)}. \quad (33)$$

so we have the collinear factorizations

$$D_{qg}^{(0)} = \Gamma_{qq}^N \left[(D_{qq} + \frac{1}{2n_f} D_{q\bar{q}}) 2n_f \Gamma_{qg} + D_{qg} \Gamma_{gg} \right], \quad (34)$$

$$(D_{qq}^{(0)} + \frac{1}{2n_f} D_{q\bar{q}}^{(0)}) = \Gamma_{qq}^N \left[(D_{qq} + \frac{1}{2n_f} D_{q\bar{q}}) \Gamma_{qq} + D_{qg} \Gamma_{gg} \right]. \quad (35)$$

Now consider first the gluonic contribution. Using eqns. (7) and eqns. (8), then since $\Gamma_{qq}^N = 1 + \mathcal{O}(\alpha_s)$, while $\Gamma_{qg} = \mathcal{O}(\alpha_s(\frac{\alpha_s}{N})^k)$

$$D_{qg}^{(0)} = \Gamma_{qg} + D_{qg} \Gamma_{gg} + \mathcal{O} \left(\alpha_s^2 \left(\frac{\alpha_s}{N} \right)^k \right). \quad (36)$$

We would like to factorize out an overall factor of Γ_{gg} , in order to compare to the k_T -factorization eqn. (12). This can be achieved by taking the logarithmic derivative with respect to Q^2 [11]: since

$$\frac{\partial}{\partial \ln Q^2} \Gamma_{gg}(N, \alpha_s(\frac{Q^2}{\mu^2})^\varepsilon, \varepsilon) = \gamma_{gg}(N, \alpha_s(\frac{Q^2}{\mu^2})^\varepsilon S_\varepsilon) \Gamma_{gg}(N, \alpha_s(\frac{Q^2}{\mu^2})^\varepsilon, \varepsilon), \quad (37)$$

$$\frac{\partial}{\partial \ln Q^2} \Gamma_{qg}(N, \alpha_s(\frac{Q^2}{\mu^2})^\varepsilon, \varepsilon) = \gamma_{qg}(N, \alpha_s(\frac{Q^2}{\mu^2})^\varepsilon S_\varepsilon) \Gamma_{qg}(N, \alpha_s(\frac{Q^2}{\mu^2})^\varepsilon, \varepsilon), \quad (38)$$

we find

$$\frac{\partial}{\partial \ln Q^2} D_{qg}^{(0)} \Big|_{\mu^2=Q^2} = \left[\gamma_{qg} + \gamma_{gg} D_{qg} + \varepsilon \alpha_s \frac{\partial}{\partial \alpha_s} D_{qg} \right] \Gamma_{gg} + \mathcal{O}(\alpha_s^2 (\frac{\alpha_s}{N})^k). \quad (39)$$

On the other hand, from eqn. (20) the high energy factorization gives

$$\frac{\partial}{\partial \ln Q^2} D_{qg}^{(0)} \Big|_{\mu^2=Q^2} = h_{qg}(N, \gamma_{gg}; \alpha_s) R(\gamma_{gg}) \Gamma_{gg}(N; \alpha_s, \varepsilon) + \mathcal{O}(\alpha_s^2 (\frac{\alpha_s}{N})^k). \quad (40)$$

Comparing eqns. (39) and (40), we can factor all the collinear singularities into Γ_{gg} on each side, and thus we must have

$$\gamma_{qg}(N, \alpha_s) + \gamma_s(\frac{\alpha_s}{N}) D_{qg}(N, \alpha_s) = h_{qg}(N, \gamma_s; \alpha_s) R(\gamma_s) + \mathcal{O}(\alpha_s^2 (\frac{\alpha_s}{N})^k), \quad (41)$$

where we also used eqn. (17). We can thus reduce the calculation of D_{qg} in $\overline{\text{MS}}$ factorization to a calculation of the impact factor eqn. (19) since γ_s , γ_{qg} and $R(\gamma_s)$ are already known to the required order [11].

The quark coefficient function D_{qg} may be determined similarly: since $D_{q\bar{q}} = 1 + \mathcal{O}(\alpha_s)$ and $D_{q\bar{q}}^{(0)} = 1 + \mathcal{O}(\alpha_s)$, eqn.(35) may be written

$$D_{qg}^{(0)} = D_{qg} + \Gamma_{qg}^{PS} + D_{qg} \Gamma_{gq} + \mathcal{O}(\alpha_s^2 (\frac{\alpha_s}{N})^k). \quad (42)$$

At high energy this is given by eqn. (21). The subtraction term is just the on-shell partonic cross-section, i.e. the bare coefficient function evaluated at $\mathcal{O}(\alpha_s)$. It may be evaluated using eqs. (36) and (29):

$$\Sigma_{qg}(N, 0; \alpha_s, \varepsilon) = \frac{\alpha_s}{\varepsilon} \gamma_{qg}^0(N) + \alpha_s D_{qg}^0(N) + \mathcal{O}(\alpha_s^2) \quad (43)$$

where the first term is the collinear singularity from the quark-gluon splitting and the second the renormalized coefficient function to $\mathcal{O}(\alpha_s)$. Substituting eqs. (36), (42) and (43) into the bare colour charge relation eqn. (21) we obtain:

$$D_{qg} + \Gamma_{qg}^{PS} + D_{qg} \Gamma_{gq} = \frac{C_F}{C_A} [\Gamma_{qg} + D_{qg} \Gamma_{gg} - \frac{\alpha_s}{\varepsilon} \gamma_{qg}^0 - \alpha_s D_{qg}^0] + \mathcal{O}(\alpha_s^2 (\frac{\alpha_s}{N})^k). \quad (44)$$

Using the colour charge relations (31) and (30) all the singular terms cancel as they should, and we are left with the renormalized colour charge relation for the Drell-Yan coefficient functions

$$D_{qg}(N, \alpha_s) = \frac{C_F}{C_A} [D_{qg}(N, \alpha_s) - \alpha_s D_{qg}^0(0)] + \mathcal{O}(\alpha_s^2 (\frac{\alpha_s}{N})^k). \quad (45)$$

Thus the high energy behaviour of the pure singlet quark coefficient function D_{qq} may be determined directly from that of the quark gluon coefficient function D_{qg} found from the eqn. (19) through eqn. (41).

Finally we consider the factorization of the remaining contribution to the Drell-Yan cross-section, σ^{SS} given by eqn. (26). The terms in the first line are $\mathcal{O}(\alpha_s(\frac{\alpha_s}{N})^k)$, while those in the second are $\mathcal{O}(\alpha_s^2(\frac{\alpha_s}{N})^k)$, so we consider them separately. From the coefficients of $Q^{(0)}G^{(0)}$ and $Q^{(0)}Q^{(0)}$ in the first line we now find (in the same way that we derived eqns. (34) and (35))

$$D_{qg}^{(0)} = 2(D_{qq} + \frac{1}{4n_f}D_{q\bar{q}})\Gamma_{qq}2n_f\Gamma_{qg} + D_{qg}(\Gamma_{qq}\Gamma_{gg} + 2n_f\Gamma_{qg}\Gamma_{gq}), \quad (46)$$

$$(D_{qq}^{(0)} + \frac{1}{4n_f}D_{q\bar{q}}^{(0)}) = (D_{qq} + \frac{1}{4n_f}D_{q\bar{q}})\Gamma_{qq}^2 + D_{qg}\Gamma_{qq}\Gamma_{gq}. \quad (47)$$

Keeping only terms of $\mathcal{O}(\alpha_s(\frac{\alpha_s}{N})^k)$, these simplify to eqns. (36) and (42), as they must.

From the second line of eqn. (26) consider the gluon-gluon term: it is easy to show that this may be used to determine the resummed coefficient function D_{gg} in terms of an impact factor computed as the double Mellin transform of the second derivative of the off-shell partonic cross-section $\Sigma_{gg}(N, \frac{\mathbf{k}_1^2}{Q^2}, \frac{\mathbf{k}_2^2}{Q^2}; \alpha_s, \varepsilon)$ in eqn. (12): the result is

$$\gamma_{qg}^2 + \gamma_{qg}\gamma_s D_{qg} + \gamma_s^2 D_{gg} = h_{gg}(N, \gamma_s, \gamma_s; \alpha_s)R(\gamma_s)^2 + \mathcal{O}(\alpha_s^2(\frac{\alpha_s}{N})^k), \quad (48)$$

where

$$h_{gg}(N, M_1, M_2; \alpha_s) \equiv M_1^2 M_2^2 \int_0^\infty d\xi_1 \xi_1^{M_1-1} \int_0^\infty d\xi_2 \xi_2^{M_2-1} \Sigma_{gg}(N, \xi_1, \xi_2; \alpha_s, 0). \quad (49)$$

However in this case the bare colour-charge relations implicit in the k_T -factorization do not lead directly to simple colour-charge relations among the coefficients $\hat{D}_{qq}$, $\hat{D}_{qg}$ and D_{gg} , as they did for heavy quark production [14], because here there are also contributions of the same order from $\mathcal{O}(\alpha_s^2(\frac{\alpha_s}{N})^k)$ terms in D_{qq} and D_{qg} , and these cannot at present be computed until k_T -factorization is extended to NNLLx.

For the remainder of this paper, we will work entirely at NLLx, computing the impact factor h_{qg} and thus all the $\mathcal{O}(\alpha_s(\frac{\alpha_s}{N})^k)$ contributions to the coefficient functions D_{qq} and D_{qg} .

5 The off-shell cross-section

In this section we compute the off-shell cross-section $\Sigma_{qg}(\tau, \xi; \alpha_s, 0)$ for the process

$$g^*(k) q(p) \rightarrow \gamma^*(q) q(p'),$$

where

$$\begin{aligned} k &= x_1 p_1 + \mathbf{k}, & p &= x_2 p_2, \\ q &= z_1 x_1 p_1 + (1 - z_2)x_2 p_2 + \mathbf{q}, \\ p' &= (1 - z_1)x_1 p_1 + z_2 x_2 p_2 + \mathbf{k} - \mathbf{q}, \end{aligned} \quad (50)$$

and we define the dimensionless variables

$$\tau = \frac{Q^2}{\nu}, \quad \xi = \frac{|\mathbf{k}|^2}{Q^2}, \quad (51)$$

with $Q^2 = q^2$ and $\nu = 2x_1x_2p_1 \cdot p_2$. We need to consider two diagrams, given in fig. 2. Since soft and collinear divergences are regulated by the off-shellness of the incoming gluon, we may perform the whole calculation in four dimensions. The two-body phase-space is then given by

$$\begin{aligned}
d\Phi^{(2)} &= \frac{d^4q}{(2\pi)^3} \frac{d^4p'}{(2\pi)^3} \delta(q^2 - Q^2) \delta(p'^2) (2\pi)^4 \delta^{(4)}(k + p - q - p') \\
&= \frac{d^4q}{(2\pi)^2} \delta(q^2 - Q^2) \delta(p'^2) = \\
&= \frac{\nu}{8\pi^2} dz_1 dz_2 d^2\mathbf{q} \delta((1 - z_2)z_1\nu - |\mathbf{q}|^2 - Q^2) \delta((1 - z_1)z_2\nu - |\mathbf{k} - \mathbf{q}|^2). \tag{52}
\end{aligned}$$

The squared matrix element is computed using the usual eikonal polarisations for the gluon; the photon indices are contracted with $g^{\mu\nu}$, because of conservation of electro-magnetic current, just as in the on-shell calculation [1]. The result is

$$\begin{aligned}
|\mathcal{M}|^2 &= -\frac{e_q^2 g_s^2}{N_c} \left\{ \frac{t}{s} + \frac{s}{t} + Q^2 |\mathbf{k}|^2 \left(\frac{1}{s^2} + \frac{1}{t^2} \right) + 4 \frac{\mathbf{k} \cdot \mathbf{q}}{s} - 4 \frac{Q^2 \mathbf{k} \cdot \mathbf{q}}{t^2} \left(1 - \frac{\mathbf{k} \cdot \mathbf{q}}{|\mathbf{k}|^2} \right) \right. \\
&\quad \left. + \frac{2}{st} \left[(Q^2 + |\mathbf{k}|^2)(|\mathbf{k}|^2 - 2\mathbf{k} \cdot \mathbf{q}) + 2(\mathbf{k} \cdot \mathbf{q})^2 + |\mathbf{k}|^2(s - t) \right] \right\} \tag{53}
\end{aligned}$$

where the Mandelstam invariants are

$$\begin{aligned}
s &= (p + k)^2 = \nu - |\mathbf{k}|^2, \\
t &= (p - q)^2 = Q^2 - z_1\nu, \\
u &= Q^2 - |\mathbf{k}|^2 - s - t. \tag{54}
\end{aligned}$$

In the on-shell limit $|\mathbf{k}| \rightarrow 0$, when averaged over the angle ϑ between $\mathbf{k}$ and $\mathbf{q}$, the off-shell matrix element eqn. (53) reduces to

$$\lim_{|\mathbf{k}| \rightarrow 0} \langle |\mathcal{M}|^2 \rangle_\vartheta = -\frac{e_q^2 g_s^2}{N_c} \left\{ \frac{t}{s} + \frac{s}{t} + 2 \frac{Q^2 u}{st} \right\}, \tag{55}$$

in agreement with the standard calculation.

To perform the phase space integration we first change the variable of integration by introducing $\Delta = \mathbf{q} - z_1 \mathbf{k}$, and we then use one delta function to perform the integral over z_2 , and the second to perform the integral over $\Delta^2 = |\mathbf{q}|^2 - 2z_1 \mathbf{k} \cdot \mathbf{q} + z_1^2 |\mathbf{k}|^2$, i.e. we write

$$\begin{aligned}
d\Phi^{(2)} &= \frac{1}{8\pi^2} \frac{dz_1 dz_2}{(1 - z_1)} d^2\Delta \delta((1 - z_2)z_1\nu - |\mathbf{q}|^2 - Q^2) \delta(z_2 - \frac{|\mathbf{k} - \mathbf{q}|^2}{(1 - z_1)\nu}) \\
&= \frac{1}{16\pi^2} dz_1 d\vartheta dz_2 d\Delta^2 \delta(z_2 - \frac{|\mathbf{k} - \mathbf{q}|^2}{(1 - z_1)\nu}) \delta(\Delta^2 - (1 - z_1)[z_1(\nu - |\mathbf{k}|^2) - Q^2]). \tag{56}
\end{aligned}$$

The remaining integral over ϑ then runs from zero to 2π , while the integral over z_1 runs from $Q^2/(\nu - |\mathbf{k}|^2)$ to one. The dimensionless cross-section can then be written as

$$\Sigma_{qg}(\tau, \xi; \alpha_s, 0) = \frac{\alpha_s}{2\pi} \frac{\tau}{2} \int_{\frac{\tau}{1-\tau\xi}}^1 dz_1 \int_0^{2\pi} \frac{d\vartheta}{2\pi} |\mathcal{M}|^2|_{z_2, \Delta^2}, \tag{57}$$

where $|\mathcal{M}|^2|_{z_2, \Delta^2}$ is the squared matrix element (53) with the two δ -functions (56) imposed. The normalization is consistent with eqn. (2), with in particular the factor $\frac{e_q^2}{N_c}$ has been absorbed into the LO cross-section. The two remaining integrations can now be performed explicitly: the result is

$$\begin{aligned} \Sigma_{qg}(\tau, \xi; \alpha_s, 0) = & \frac{\alpha_s}{2\pi} T_R \tau \left\{ \frac{1}{1 - \tau\xi} \ln \left(\frac{(1-\tau)(1-\xi\tau)}{\tau^2\xi} \right) p_1(\tau, \xi) \right. \\ & \left. + \frac{1 - \tau - \tau\xi}{2(1 - \tau)(1 - \tau\xi)^3} p_2(\tau, \xi) \right\} \Theta \left(\frac{1}{\tau} - \xi - 1 \right). \end{aligned} \quad (58)$$

where the polynomials

$$\begin{aligned} p_1(\tau, \xi) &= \tau^2 + (1 - \tau)^2 + \tau^2\xi (10 + \xi + 18\tau^2\xi - 6\tau(3 + 2\xi)), \\ p_2(\tau, \xi) &= -1 + 36\tau^5\xi^3 + 7\tau(2 + \xi) - 6\tau^4\xi^2(15 + 7\xi) \\ &\quad + 2\tau^3\xi(35 + 49\xi + 4\xi^2) - \tau^2(15 + 71\xi + 14\xi^2). \end{aligned} \quad (59)$$

Note that this cross-section has been already averaged over the azimuthal angle of the incoming gluon, consistently with eqns. (18) and (19), where the Mellin integral is taken with respect to $\xi = \mathbf{k}^2/Q^2$. The on-shell limit $\xi \rightarrow 0$ is

$$\Sigma_{qg}(\tau, \xi; \alpha_s, 0) = \frac{\alpha_s}{2\pi} \left[\tau P_{qg}(\tau) \ln \left(\frac{1-\tau}{\xi\tau^2} \right) + \frac{1}{2}\tau \left(-\frac{1}{2} + 7\tau - \frac{15}{2}\tau^2 \right) \right] + \mathcal{O}(\xi), \quad (60)$$

where $P_{qg}(\tau) = T_R(\tau^2 + (1 - \tau)^2)$ is the LO qg -splitting function. Because of the off-shell regularisation, this result is not the same as the usual $\overline{\text{MS}}$ NLO coefficient function [1]:

$$D_{qg}^1(\tau; \alpha_s, \varepsilon) = \frac{\alpha_s}{2\pi} \left[\tau P_{qg}(\tau) \left(\frac{1}{\varepsilon} + \ln 4\pi - \gamma_E + \ln \left(\frac{(1-\tau)^2}{\tau} \right) \right) + \frac{1}{2}\tau \left(\frac{1}{2} + 3\tau - \frac{7}{2}\tau^2 \right) \right], \quad (61)$$

though the collinear singularities match as they must, with $\ln(1/\xi) \rightarrow 1/\varepsilon + \ln 4\pi - \gamma_E$.

6 The quark-gluon impact factor

The quark-gluon impact factor is defined as the double Mellin transform of the double logarithmic derivative of the off-shell cross-section (see eqn. (19):

$$h_{qg}(N, M; \alpha_s) = M^2 \int_0^\infty \xi^{M-1} \int_0^1 \tau^{N-1} \Sigma(\tau, \xi; \alpha_s, 0). \quad (62)$$

Because the off-shell cross-section eqn. (58) contains a factor $\Theta(\tau^{-1} - \xi - 1)$, it is useful to change the integration variables according to

$$\alpha = \tau\xi, \quad \beta = \frac{\tau}{1 - \tau\xi}; \quad (63)$$

the Jacobian determinant is $1/\beta$ and the Θ -function condition is satisfied for all $\alpha, \beta \in [0, 1]$, so

$$h_{qg}(N, M; \alpha_s, 0) = \frac{\alpha_s}{2\pi} T_R M^2 \int_0^1 d\alpha \int_0^1 d\beta \alpha^{M-1} (1-\alpha)^{N-M} \beta^{N-M} \left[\ln \frac{1-(1-\alpha)\beta}{\alpha\beta} d_1(\alpha, \beta) - \frac{1}{2} \frac{(1-\beta)}{1-(1-\alpha)\beta} d_2(\alpha, \beta) \right], \quad (64)$$

where the polynomials d_1 and d_2 are now

$$\begin{aligned} d_1(\alpha, \beta) &= 1 + \alpha^2 - 2\beta + 12\alpha\beta - 22\alpha^2\beta + 12\alpha^3\beta + 2\beta^2 - 22\alpha\beta^2 \\ &\quad + 56\alpha^2\beta^2 - 54\alpha^3\beta^2 + 18\alpha^4\beta^2, \\ d_2(\alpha, \beta) &= 1 - 6\alpha + 8\alpha^2 - 14\beta + 71\alpha\beta - 98\alpha^2\beta + 42\alpha^3\beta + 15\beta^2 \\ &\quad - 85\alpha\beta^2 + 160\alpha^2\beta^2 - 126\alpha^3\beta^2 + 36\alpha^4\beta^2. \end{aligned} \quad (65)$$

The integrals containing the logarithm can be further simplified through integrating by parts with respect to the variable β :

$$\begin{aligned} &\int_0^1 d\alpha \int_0^1 d\beta \alpha^{M-1+p} (1-\alpha)^{N-M} \beta^{N-M+q} \ln \frac{1-(1-\alpha)\beta}{\alpha\beta} = \\ &\frac{1}{1+N-M+q} \int_0^1 d\alpha \int_0^1 d\beta \alpha^{M-1+p} (1-\alpha)^{N-M} \beta^{N-M+q} \frac{1}{1-(1-\alpha)\beta}. \end{aligned} \quad (66)$$

where p and q are integers, the exponents of α and β in each terms of the functions $d_1(\alpha, \beta)$. Thus the only nontrivial integral which we need is

$$\begin{aligned} &\int_0^1 d\alpha \int_0^1 d\beta \alpha^{M-1+p} (1-\alpha)^{N-M} \beta^{N-M+q} \frac{1}{1-(1-\alpha)\beta} = \\ &= \sum_{k=0}^{\infty} \int_0^1 d\alpha \int_0^1 d\beta \alpha^{M-1+p+k} (1-\alpha)^{N-M} \beta^{N-M+q+k} \\ &= \sum_{k=0}^{\infty} \frac{\Gamma(1+N-M+k)\Gamma(M+p)}{\Gamma(1+N+k+p)(1+N-M+q+k)} \end{aligned} \quad (67)$$

$$\begin{aligned} &= \frac{\Gamma(M+p)\Gamma(1+N-M)}{\Gamma(1+N+p)(1+N-M+q)} \\ &\quad {}_3F_2(1, 1+N-M, 1+N-M+q; 1+N+p, 2+N-M+q; 1) \\ &= \Gamma^2(M+p) \frac{\Gamma(1+N-M)\Gamma(1+N-M+q)}{\Gamma(1+N+p)\Gamma(1+N+p+q)} \\ &\quad {}_3F_2(M+p, N+p, 1+N-M+q; 1+N+p, 1+N+p+q; 1), \end{aligned} \quad (68)$$

where ${}_3F_2(a, b, c; d, e; z)$ is a generalised hypergeometric function, and in the last line we used Thomae's theorem. The final result for h_{qg} eqn. (64) is then a fairly complicated sum of terms involving the

generalised hypergeometric functions ${}_3F_2$:

$$\begin{aligned}
h_{qg}(N, M; \alpha_s, 0) &= \frac{\alpha_s}{2\pi} T_R \sum_{p=0}^4 \sum_{q=0}^3 M^2 \Gamma^2(M+p) \frac{\Gamma(1+N-M)\Gamma(1+N-M+q)}{\Gamma(1+N+p)\Gamma(1+N+p+q)} \\
&\quad {}_3F_2(M+p, N+p, 1+N-M+q; 1+N+p, 1+N+p+q; 1) \\
&\quad \left[\frac{\Delta_{1+p,1+q}^{(1)}}{(1+N-M+q)} + \Delta_{1+p,1+q}^{(2)} \right]
\end{aligned} \tag{69}$$

where

$$\Delta_{i,j}^{(1)} = \begin{pmatrix} 1 & -2 & 2 & 0 \\ 0 & 12 & -22 & 0 \\ 1 & -22 & 56 & 0 \\ 0 & 12 & -54 & 0 \\ 0 & 0 & 18 & 0 \end{pmatrix}, \quad \Delta_{i,j}^{(2)} = -\frac{1}{2} \begin{pmatrix} 1 & -15 & 29 & -15 \\ -6 & 77 & -156 & 85 \\ 8 & -106 & 258 & -160 \\ 0 & 42 & -168 & 126 \\ 0 & 0 & 36 & -36 \end{pmatrix}. \tag{70}$$

Note that in the collinear limit $M \rightarrow 0$ eqn. (69) is regular. In fact since only terms with $p = 0$ survive, and ${}_3F_2(0, N, 1+N+q; 1+N, 1+N+q; 1) = 1$,

$$h_{qg}(N, 0; \alpha_s, 0) = \frac{\alpha_s}{2\pi} T_R \frac{N^2 + 3N + 4}{(N+1)(N+2)(N+3)} = \alpha_s \gamma_{qg}^0(N) \tag{71}$$

as it must be from eqn. (41). This is a highly nontrivial check on eqn. (69).

In the limit $N \rightarrow 0$ (i.e. in the high energy limit), the sum (67) may be performed more easily, since the generalised hypergeometric function reduces to a sum of rational functions of Γ -functions:

$$\begin{aligned}
&\int_0^1 d\alpha \int_0^1 d\beta \alpha^{M-1+p} (1-\alpha)^{-M} \beta^{-M+q} \frac{1}{1-(1-\alpha)\beta} \\
&= \sum_{k=0}^{\infty} \frac{\Gamma(1-M+k)\Gamma(M+p)}{\Gamma(1+k+p)(1-M+q+k)} \\
&= \Gamma(M+p) \left[\frac{\Gamma(M+p)\Gamma(1-M-p)\Gamma(1-M-p+q)}{\Gamma(1+q)} \right. \\
&\quad \left. - \sum_{j=0}^{p-1} \frac{\Gamma(1-M-p+j)}{(1-M+q-p+j)j!} \right].
\end{aligned} \tag{72}$$

Using this in eqn. (64), the result remarkably simplifies to

$$h_{qg}(0, M; \alpha_s, 0) = \frac{\alpha_s}{2\pi} T_R 4 \frac{\Gamma(1-M)^2 \Gamma(1+M)^2}{(1-M)(2-M)(3-M)}. \tag{73}$$

The same result may be obtained by setting $N = 0$ in the generalised hypergeometric functions in eqn. (69), and simplifying the result. The infrared singularity at $M = 1$ is a triple pole, as expected from the counting of soft and collinear singularities [12].

7 Drell-Yan coefficient functions at high energy

The Taylor expansion of eqn. (73) about $M = 0$ gives

$$h_{qg}(0, M; \alpha_s, 0) = \frac{\alpha_s}{2\pi} T_R \frac{2}{3} \left[1 + \frac{11}{6} M + \left(\frac{85}{36} + \frac{\pi^2}{3} \right) M^2 + \left(\frac{575}{216} + \frac{11\pi^2}{18} \right) M^3 + \left(\frac{3661}{1296} + \frac{85\pi^2}{108} + \frac{\pi^4}{15} \right) M^4 + \mathcal{O}(M^5) \right] \quad (74)$$

In order to compute the high energy behaviour of the $\overline{\text{MS}}$ coefficient function, we need the scheme dependent factor:

$$R(M) = 1 + \frac{8}{3}\zeta_3 M^3 - \frac{3}{4}\zeta_4 M^4 + \mathcal{O}(M^5). \quad (75)$$

The BFKL anomalous dimension is

$$\gamma_s \left(\frac{\alpha_s}{N} \right) = \frac{C_A \alpha_s}{\pi N} + 2\zeta_3 \left(\frac{C_A \alpha_s}{\pi N} \right)^4 + 2\zeta_5 \left(\frac{C_A \alpha_s}{\pi N} \right)^6 + \dots \quad (76)$$

while quark anomalous dimension in $\overline{\text{MS}}$ may be found in [11]:

$$\begin{aligned} \gamma_{qg}(N, \alpha_s) &= \frac{\alpha_s}{3\pi} T_R \left[1 + \frac{5}{3} \frac{C_A}{\pi} \frac{\alpha_s}{N} + \frac{14}{9} \left(\frac{C_A}{\pi} \frac{\alpha_s}{N} \right)^2 + \left(\frac{82}{81} + 2\zeta_3 \right) \left(\frac{C_A}{\pi} \frac{\alpha_s}{N} \right)^3 \right. \\ &\quad \left. + \left(\frac{122}{243} + \frac{25}{6}\zeta_3 \right) \left(\frac{C_A}{\pi} \frac{\alpha_s}{N} \right)^4 + \left(\frac{146}{729} + \frac{14}{3}\zeta_3 + 2\zeta_5 \right) \left(\frac{C_A}{\pi} \frac{\alpha_s}{N} \right)^5 + \dots \right]. \end{aligned} \quad (77)$$

Substituting eqn. (76) into eqns. (74) and (75), and then substituting the results along with eqn. (77) into eqn. (41), the leading terms on either side cancel, and the remaining terms give the NLL x expansion

$$\begin{aligned} D_{qg}(N, \alpha_s) &= \frac{\alpha_s}{18\pi} T_R \left[1 + \left(\frac{29}{6} + 2\pi^2 \right) \frac{C_A}{\pi} \frac{\alpha_s}{N} \right. \\ &\quad \left. + \left(\frac{1069}{108} + \frac{11}{3}\pi^2 + 4\zeta_3 \right) \left(\frac{C_A}{\pi} \frac{\alpha_s}{N} \right)^2 \right. \\ &\quad \left. + \left(\frac{9031}{648} + \frac{85}{18}\pi^2 + \frac{7}{20}\pi^4 + \frac{73}{3}\zeta_3 \right) \left(\frac{C_A}{\pi} \frac{\alpha_s}{N} \right)^3 + \dots \right]. \end{aligned} \quad (78)$$

The coefficients of $\mathcal{O}(\alpha_s)$ and $\mathcal{O}(\alpha_s^2)$ are in agreement with the high energy limit of the fixed order NLO [1] and NNLO [3, 4] computations. This is a very non-trivial check of the procedure. The $\mathcal{O}(\alpha_s^3)$ and subsequent terms are all new results.

The high energy singularities of the quark-quark coefficient function are now easily deduced using

the colour-charge relation eqn. (45): we find

$$\begin{aligned}
D_{qq}(N, \alpha_s) = & \frac{\alpha_s}{18\pi} T_R \frac{C_F}{\pi} \frac{\alpha_s}{N} \left[\left(\frac{29}{6} + 2\pi^2 \right) \right. \\
& + \left(\frac{1069}{108} + \frac{11}{3}\pi^2 + 4\zeta_3 \right) \frac{C_A}{\pi} \frac{\alpha_s}{N} \\
& \left. + \left(\frac{9031}{648} + \frac{85}{18}\pi^2 + \frac{7}{20}\pi^4 + \frac{73}{3}\zeta_3 \right) \left(\frac{C_A}{\pi} \frac{\alpha_s}{N} \right)^2 + \dots \right]. \quad (79)
\end{aligned}$$

Again this result checks at $\mathcal{O}(\alpha_s^2)$ against the NNLO result of ref. [3, 4], while the terms from $\mathcal{O}(\alpha_s^3)$ onwards are new results.

8 Summary

We have analysed the Drell-Yan (and thus also vector boson production) process in the limit of high partonic centre-of-mass energy, along the lines of the analysis of DIS presented in [11]. We showed how to factorise simultaneously high energy and collinear singularities from the hard coefficient function, computed the quark-gluon impact factor, and thus deduced the leading high energy singularities in the Drell-Yan coefficient functions in $\overline{\text{MS}}$ scheme to arbitrarily high orders in α_s . Our results agree with the known results at NLO and NNLO, while providing new results at N³LO and beyond.

It will now be possible to use these results to perform an all order resummation of high energy logarithms in Drell-Yan processes, using the techniques developed in [12] which have already been applied to deep inelastic processes [13]. A global analysis of the effect of high energy resummation on parton distribution functions and LHC benchmarks is thus now a definite possibility for the near future.

Acknowledgements: SM would like to thank SUPA for financial support during most of the period during which this work was done. We were also supported in part by the Marie Curie Research and Training network HEPTOOLS under contract MRTN-CT-2006-035505.

References

- [1] G. Altarelli, R. K. Ellis and G. Martinelli, Nucl. Phys. B **143** (1978) 521 [Erratum-ibid. B **146** (1978) 544].
- [2] T. Matsuura, R. Hamberg and W. L. van Neerven, Nucl. Phys. B **345** (1990) 331.
- [3] R. Hamberg, W. L. van Neerven and T. Matsuura, Nucl. Phys. B **359** (1991) 343 [Erratum-ibid. B **644** (2002) 403].
- [4] J. Blumlein and V. Ravindran, Nucl. Phys. B **716**, 128 (2005), [hep-ph/0501178].
- [5] S. Catani and L. Trentadue, Nucl. Phys. B **327** (1989) 323.
- [6] S. Catani, D. de Florian, M. Grazzini and P. Nason, JHEP **0307**, 028 (2003), [arXiv:hep-ph/0306211].
- [7] S. Moch and A. Vogt, Phys. Lett. B **631** (2005) 48, [hep-ph/0508265].
- [8] E. Laenen and L. Magnea, Phys. Lett. B **632** (2006) 270, [hep-ph/0508284].
- [9] G. Altarelli, R. D. Ball and S. Forte, PoS **RADCOR2007** (2007) 028 [arXiv:0802.0968 [hep-ph]].
- [10] S. Catani, M. Ciafaloni and F. Hautmann, Phys. Lett. B **242** (1990) 97; S. Catani, M. Ciafaloni and F. Hautmann, Nucl. Phys. B **366** (1991) 135.
- [11] S. Catani and F. Hautmann, Nucl. Phys. B **427** (1994) 475.
- [12] R. D. Ball, Nucl. Phys. B **796** (2008) 137, [arXiv:0708.1277 [hep-ph]].
- [13] G. Altarelli, R. D. Ball and S. Forte, Nucl. Phys. B **799** (2008) 199, [arXiv:0802.0032 [hep-ph]].
- [14] R. D. Ball and R. K. Ellis, JHEP **0105** (2001) 053, [hep-ph/0101199].
- [15] G. Camici and M. Ciafaloni, Nucl. Phys. B **496** (1997) 305 [Erratum-ibid. B **607** (2001) 431].
- [16] F. Hautmann, Phys. Lett. B **535** (2002) 159, [arXiv:hep-ph/0203140].
- [17] S. Marzani, R. D. Ball, V. Del Duca, S. Forte and A. Vicini, Nucl. Phys. B **800** (2008) 127, [arXiv:0801.2544 [hep-ph]].
- [18] S. Marzani, R. D. Ball, V. Del Duca, S. Forte and A. Vicini, Nucl. Phys. Proc. Suppl. **186** (2009) 98 [arXiv:0809.4934 [hep-ph]].